

**TITLE OF THE INVENTION**

**COMPOSITIONS AND METHODS FOR TREATMENT  
OF STAPHYLOCOCCAL INFECTION WHILE SUPPRESSING  
FORMATION OF ANTIBIOTIC-RESISTANT STRAINS**

5      *INS A1*

**BACKGROUND OF THE INVENTION**

**Field of the Invention**

006460 27099990  
This invention pertains to a method of treating staphylococcal infection in mammals, including humans. The method involves the simultaneous administration of a lysostaphin or other agent which attacks the glycine-containing peptide cross-links of the cell wall peptidoglycan found in staphylococci and an antibiotic, the antibiotic properties of which are mediated by its ability to affect the cell wall of the target staphylococci. This combined administration is effective in treating the staphylococcal infection, and at the same time suppresses the formation of strains resistant to lysostaphin or other peptidoglycan active agent.

15      **BACKGROUND OF THE PRIOR ART**

Lysostaphin is a bacteriocin secreted by a staphylococcal strain isolated and originally named *Staphylococcus staphylolyticus* (now *S. simulans*). The production of lysostaphin is described in U.S. Patent 3,278,378. Lysostaphin is an endopeptidase which cleaves the polyglycine cross-links of the peptidoglycan found in the cell walls of staphylococci. U.S. Patents 3,398,056 and 3,594,284 describe improvements to culture medium and inoculation techniques for the production of lysostaphin.

20

10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995

The gene for lysostaphin from *S. simulans* has been sequenced and cloned, U.S. Patent 4,931,390. Lysostaphin for use as a laboratory reagent has been produced by fermentation of a non-pathogenic recombinant strain of *B. sphaericus*, from which it is readily purified. The cloning and sequencing of the lysostaphin gene permits the isolation of variant enzymes that have properties similar to or different from those of wild type lysostaphin. One such altered enzyme, bearing a single amino acid change, has been characterized and shown to have potent anti-staphylococcal activity both *in vitro* and in an animal infection model. U.S. Patent Application Serial No. 09/120,030, filed July 21, 1998 and incorporated herein by reference. Other lysostaphin analogues, including naturally occurring enzymes of this type have been established as potent agents capable of addressing difficult to treat bacterial diseases caused by staphylococcal infection. Other peptidases with related activity are known. Thus *lasA* protease and achromopeptidase, reported in Kessler, et al., J. Biol. Chem. 268:7503-08 (1993) and Li et al., J. Biochem. 122:772-778(1997), respectively, have anti-staphylococcal activity based on their digestion of glycine-containing cross-links in the peptidoglycan cell wall component. These agents may be used in this invention in place of lysostaphin.

The development of lysostaphin as an effective antibiotic to treat staphylococcal infection has been plagued, however, by a problem that is universal for antibiotic administration - the increasing development of antibiotic-resistant strains of mutant staphylococci. Already, a wide variety of staphylococcal infections resistant to various antibiotics that were previously the treatment of choice, including methicillin (methicillin resistant *S. aureus* are referred to as MRSA) and vancomycin-resistant strains (referred to as VISA) have been identified. Resistance to a wide variety of other antibiotics, not

exhibited by sensitive staphylococci, has been noted as well. MRSA, as well as strains resistant to other antibiotics, are discussed at length in Stranden, et al., J. Bacteriology 179(1):9-16 (1997). Further difficulties are encountered in that MRSA tend to accumulate a variety of other resistances as well. Multiresistant MRSA are typically treated with vancomycin, The Staphylococci In Human Diseases, 158-174 (Grossley, et al., editors 5 1997). Vancomycin itself may be toxic. Additionally, vancomycin resistance has recently been detected in staphylococci infections.

The problem posed by the continuing development of antibiotic-resistant infectious agents, such as staphylococci, is more than the difficulty involved in treating any individual patient. Popular press, as well as scientific journals, have noted the alarming increase in the generation of resistant strains, due in part to indiscriminate use or over-use of antibiotics. Each time an individual is treated with an antibiotic, whether needlessly or reasonably, the chance that a strain resistant to that particular treatment will arise is increased. Resistant strains of staphylococci have become endemic in many hospitals and pose a life-threatening danger to patients already debilitated by other ailments who become infected after admission to those hospitals.

10  
15  
20  
SUB A27  
Numerous articles have noted the development of resistance to either lysostaphin or  $\beta$ -lactams, such as methicillin, and the relationship there between. Thus, DeHart, et al., Applied Environmental Microbiology 61, 1475-1479 (1995) noted the development of mutant *S. aureus* recombinant cells that were resistant to lysostaphin, but susceptible to methicillin. Similar phenomenon are reported by Zygmunt, et al., Can. J. Microbio. 13,845-852 (1966), Polak, et al., Diagn. Microbiol. Infet. Dis. 17:265-270 (1993) and Dickson, et al., Yale J. Bio. Med. 41:62-67 (1968). Each of these references, as well as

later reports such as Ehlert, J. Bacteriology, 179:7573-7576 (1997) note that staphylococci  
SUB AZ  
cont that develop resistance to lysostaphin, either spontaneously or through induced  
recombination, become susceptible to methicillin treatment, and vice-versa. In all of these  
references, the uniform suggestion is to follow a course of administration of lysostaphin,  
5 even a short one, with administration of methicillin.

U.S. Patent 5,760,026, commonly assigned herewith, employs a specific method for  
treating mastitis, by intramammary infusion of lysostaphin. The patent reports, Table ID  
and elsewhere, that a synergistic result is predicted when combining lysostaphin and a  $\beta$ -  
lactam to treat mastitis, based on an *in vitro* assay. The bovine mastitis model is not  
predictive of *in vivo* administration of antibiotics, and the synergistic effects reported in  
U.S. Patent 5,760,026 have not been substantiated in an environment or model that would  
be reflective of *in vivo* administration to a mammal such as a human.

Those of skill in the art will be aware that there are a wide variety of staphylococcal  
strains. Many are resistant to conventional antibiotics, unlike sensitive strains. *S. aureus*  
strains are recognized as highly virulent and the most common single cause of serious  
systemic infections. Coagulase-negative staphylococcal species, although generally less  
invasive than *S. aureus*, are now responsible for a significant incidence of infections;  
particular among debilitated or immunocompromised patients. As an example of such  
infection, one may point to endocarditis consequent to heart valve replacement. This is but  
20 one of a variety of intractable staphylococcal infections which are increasing due to the  
widespread use of antibiotics.

Accordingly, it remains an object of those of ordinary skill in the art to develop a  
method whereby even resistant staphylococcal infections in mammals, including humans,

may be effectively treated by the administration of antibiotics. Desirably, this method is developed so as suppress the formation of strains resistant to the antibiotics used.

## SUMMARY OF THE INVENTION

5 The above goals, and others made clear by the discussions set forth below, are achieved by the simultaneous administration of an anti-staphylococcal agent, such as lysostaphin or other agent whose activity is mediated by cleavage of glycine-containing cross-links in the staphylococcal cell wall peptidoglycan and an antibiotic or antimicrobial agent whose activity is mediated by its ability to affect the cell wall of staphylococci. These cell-wall active agents include  $\beta$ -lactams and glycopeptides. Preferably, the cell-wall active antibiotic is a  $\beta$ -lactam.

There is no evidence of any synergistic effect achieved through the simultaneous administration of an anti-staphylococcal agent whose activity is mediated by cleavage of glycine-containing cross-links and a cell-wall active antibiotic in a model, *in vitro* or *in vivo*, that is predictive of benefit for *in vivo* administration of antibiotics in a mammal.

15 Indeed, those of ordinary skill in the art will recognize that for resistant staphylococci, such as MRSA, the administration of methicillin is not therapeutically effective in any amount. Surprisingly, Applicants have discovered that the combined administration of an anti-staphylococcal agent whose activity is mediated by cleavage of glycine-containing cross-links such as lysostaphin and the cell-wall active antibiotic not only effectively treats the  
20 infection, but suppresses the formation of staphylococci having resistance to the anti-staphylococcal agent whose activity is mediated by cleavage of glycine-containing cross-links.

While Applicants do not wish to be bound by this explanation, it appears that the spontaneous mutation commonly effective in conferring lysostaphin resistance in staphylococci renders the same highly susceptible to a cell-wall active antibiotic, such as methicillin. This is true even where the organism starts out as methicillin resistant.

5 Simultaneous administration of both appears to be uniformly effective in simultaneously eradicating the infection and suppressing the generation of new resistant strains. Specifically, anti-staphylococcal agents like lysostaphin cleave glycine-containing cross-links. The mutation conferring resistance to this attack renders previously resistant strains sensitive to cell wall active antibiotics.

#### DETAILED DESCRIPTION OF THE INVENTION

This invention involves the administration of a pharmaceutical composition effective in the treatment of staphylococcal infection, which composition comprises at least two active agents, one an agent like lysostaphin which cleaves the glycine-containing cross-links of the cell wall peptidoglycans of staphylococci, the other a cell-wall active antibiotic. By  
15 lysostaphin it is intended to refer herein to any enzyme, including lysostaphin wild type, a mutant or variant, or any recombinant or related enzyme that retains proteolytic activity against glycine-containing cross-links in the cell wall peptidoglycan of staphylococci. Variants may be generated by post-translational processing of the protein(either by enzymes present in a producer strain or by means of enzymes or reagents  
20 introduced at any stage of the process) or by mutation of the structural gene. Mutations may include site-deletion, insertion, domain removal and replacement mutations. They may be recombinantly expressed, or otherwise. Other anti-staphylococcal active agents

acting by cleavage of the glycine-containing peptidoglycan cross-links include *lasA* protease and achromopeptidase. Such anti-staphylococcal agents which affect the peptidoglycan cross-links are embraced by the invention, but exemplified herein by reference to lysostaphin.

5           Cell-wall active antibiotics include  $\beta$ -lactams and glycopeptides.  $\beta$ -lactams are preferred. Suitable  $\beta$ -lactams include, but are not limited to, penicillins, such as penicillin, nafcillin, oxacillin, methicillin, amoxicillin and cloxacillin. Other  $\beta$ -lactams include cephalosporins and carbapenems. Representative cephalosporins include cephalothin, cefazolin, cefamandole, ceftazidime and others. Suitable carbapenems include imepenem and meropenem.

Suitable glycopeptides include vancomycin, teicoplanin and ramoplanin.

These two agents can be combined with further agents, adjuvants and the like, but are effectively administered in a pharmaceutically acceptable carrier. Administration is typically systemic, and may be intravenous (IV), intramuscular (IM), subcutaneous (SC), intraperitoneal (IP), intrathecal or topical. No synergistic effect of combining lysostaphin and a  $\beta$ -lactam or glycopeptide or cell-wall active antibiotic has been noted in a model predictive of *in vivo* mammalian administration. Accordingly, each agent of the effective combination must be administered in a therapeutically effective amount. It is to be noted, in this regard, that the amount to be administered is that which is therapeutically effective  
20   when the lysostaphin and cell-wall active agent are administered together. Those of skill in the art will of course recognize that there is no therapeutically effective amount for, e.g., methicillin if the staphylococcal infection is an MRSA infection. Nonetheless, administration of therapeutic amounts of methicillin as determined against non-MRSA,

combined with an amount of lysostaphin effective against staphylococci that are not lysostaphin-resistant will effectively treat staphylococcal infections even where the infection is resistant to one or other antibiotic. Accordingly, applicants have referred herein to "therapeutically effective amounts" to mean amounts effective to therapeutically treat sensitive *S. aureus* infection. This simultaneous administration, as opposed to sequential administration typified by the prior art, also surprisingly results in the suppression of strains resistant against either antibiotic, or their combination.

Any of a wide variety of pharmaceutically acceptable carriers and diluents, typically buffered, may be used. Appropriate pharmaceutical carriers are known to those of skill in the art. The formulations of this invention comprise a therapeutic amount of lysostaphin and a therapeutic amount of a cell-wall active antibiotic, such that when co-administered, the staphylococcal infection, either *S. aureus* or coagulase negative staphylococci, is treated, while the generation of resistant strains is suppressed. Other active agents that do not interfere with the activity of the two antibiotics may be co-administered.

Therapeutic values will range substantially given the nature of the staphylococcal infection, the individual, and the antibiotic being used in conjunction with lysostaphin. Representative values for anti-staphylococcal active agents such as lysostaphin, range from approximately 15-150 mg/kg body weight/day for human administration, with a preferred range of 25-100 mg/kg/day. Values for  $\beta$ -lactams based on nafcillin range from 50-250 mg/kg/day, with a preferred range of 100-200 mg/kg/day and glycopeptides like vancomycin are administered over a range of 10-75 mg/kg/day, with a preferred range of 15-50 mg/kg/day.

The administration course is not substantially different from that currently administered in single antibiotic treatments, and can range from 7-28 days, although typically, courses of 7-21 days are employed, and effective in treating a wide variety of staphylococcal infections.

## EXAMPLES

To compare the development of resistant strains, growth curves for three methicillin resistant staphylococcal strains were obtained for *in vitro* growth in Mueller Hinton Broth.

Growth curves were completed in Mueller Hinton Broth (50 ml) in glass erlenmeyer flasks. Flasks were inoculated with 100  $\mu$ l of an overnight growth adjusted to 0.5 Macfarland to yield a starting concentration of approximately  $10^5$  -  $10^6$  CFU/ml. Growth curves were done in the presence of lysostaphin, lysostaphin and oxacillin (1  $\mu$ g/ml) or no antibiotics (controls). Absorbance at OD 600 was recorded at 0, 2, 4, 6 and 24 hours. At 24 hours flasks were plated on MHA, MHA with lysostaphin (6  $\mu$ g/ml) and MHA with oxacillin (6  $\mu$ g/ml) in order to screen for resistant mutants. Three methicillin resistant *Staphylococcus aureus* strains were tested: 272855, 450M and Mu3.

Growth following 24 hour incubation with lysostaphin (0.0625  $\mu$ g/ml), lysostaphin (0.0625  $\mu$ g/ml) and oxacillin (1  $\mu$ g/ml), and no antibiotics, was recorded.

The data generated led to the following conclusions:

1. The addition of oxacillin to lysostaphin led to significant suppression of growth for all three strains.
2. The presence of oxacillin suppressed the expression of lysostaphin resistance among all three strains.

In order to demonstrate the effectiveness of the claimed invention, certain experiments were conducted. Checkerboard susceptibility testing was conducted to determine whether simultaneous administration of lysostaphin and oxacillin (a  $\beta$ -lactam) would be effective in suppressing the development of resistance. Oxacillin concentrations varied between 0.0156  $\mu$ g/ml and 1  $\mu$ g/ml. Lysostaphin concentrations varied between 0.00048 and 0.9  $\mu$ g/ml. Four strains were tested for evidence of synergy between lysostaphin and oxacillin; 27619, Col, 27227 and VA348. There was no evidence of synergy or antagonism over the concentration range tested. The MIC of lysostaphin was unchanged in the presence of oxacillin in concentrations up to 1  $\mu$ g/ml for all strains tested. The overnight growth of strains in the presence of lysostaphin and oxacillin was examined. Four strains were grown overnight in drug free media (MHB), MHB with 0.1  $\mu$ g/ml of lysostaphin, MHB with lysostaphin 0.1  $\mu$ g/ml and oxacillin 1  $\mu$ g/ml, and MHB with oxacillin 1  $\mu$ g/ml. The four strains tested included 450M, Col, and their lysostaphin resistant mutants 450 M lyso and Col lyso. The results are reflected in Table 1.

TABLE 1

Isolates	Growth in the presence of			
	MHB	lysostaphin 0.1 $\mu$ g/ml	oxacillin $\mu$ g/ml	Lyso + oxacillin
450M	+	+	(+)	-
450M lyso	+	+	-	-
Col	+	+	(+)	-
Col lyso	+	+	-	-

Sub A3 } The same unpredicted result has been demonstrated through *in vivo* experiments based on the widely accepted rabbit model of aortic valve endocarditis, predictive of *in vivo*

administration to humans. When administered to staphylococcal infected rabbits at low doses (1 mg/kg bid, as compared with a minimum value of 5 mg/kg tid for sterilization) lysostaphin, as representative of anti-staphylococcal agents acting by cleavage of the glycine-containing cross-links, resulted in recovery of a number of resistant colonies, with high counts in vegetations and kidneys, while the same dosage together with nafcillin (a  $\beta$ -lactam) gave sterile kidneys, some sterile vegetations, and no resistant strains recovered. The simultaneous treatment of staphylococcal infection with suppression of resistant strain formation is an exciting and widely useful invention nowhere predicted in the art. This invention offers the possibility of treating staphylococcal infections while suppressing the generation of strains resistant to any or all active agents administered.

The inventive compositions and methods of this application have been disclosed generically, and by reference to specific example, examples are not intended to be limiting unless so indicated, and variations will occur to those of ordinary skill in the art without the exercise of inventive faculty. In particular, variations in the identity of the cell-wall active antibiotic to be co-administered with an anti-staphylococcal agent acting by cleavage of the glycine-containing cross-links, as well as various recombinant and mutant variants thereof, carriers and concentrations will occur to those of skill in the art without the exercise of inventive faculty, and remain within the scope of the invention, unless specifically excluded by the claims set forth below.